
Container System

Only fo vdered [Use A 1 ug Vial 3

MINI-I

directic

2



Rx only.

To ensure safe and proper use of this device, refer to the complete instructions for use.

Assembly

20 mm Closures

Remove vial cover (1)

- Disinfect stopper
- Peel off foil cover 2
 - Inspect adaptor for moisture Discard if found
- (3) Place vial upright
 - Hold firmly
 - Push adaptor down until vial snaps in place
 - DO NOT TWIST
 - Pull vial to ensure fully seated

Reconstitution

- (4) Squeeze bag and check vial
 - · Use only if vial fully seated and dry
 - · Bend up then down to break seal
- (5) Hold bag with vial down
 - Squeeze solution into vial until half full
 - · Shake to suspend drug in solution
- (6) Hold bag with vial upside down
 - · Squeeze bag to force air into vial
 - Release to drain suspended drug from vial
 - Repeat steps 5 and 6 until vial is **empty** of drug and solution is thoroughly mixed. Ensure drug is completely dissolved. Do Not Remove Drug Vial.
- (7) Remove port protector.
 - · Attach administration set per its directions.
- (8) Hang container on I.V. pole and prime set per directions. Ensure that vial is empty of drug and solution. Repeat step 6 if drug and solution remain in vial.
 - Warning: Do not use in series connections
- (9) Administer medication per directions. Use within specified time for drug stability. Refer to drug package insert.

o Order Baxter Products, call 1-888-229-0001

Options at work.

MEDICATION DELIVERY



11094 5K 04/09

Baxte Baxte

сомм